

CompHEP: developments and applications 2016

Speaker: Mikhail Dubinin

SINP, Moscow State University

CompHEP Collaboration:

E. Boos, V. Bunichev, M. D., L. Dudko, V. Ilyin, A. Kryukov, V. Edneral, V. Savrin, A. Semenov, A. Sherstnev

<http://comphep.sinp.msu.ru>

Outline

- **History and statistics**
- **Tools for BSM physics: combined global fits, operations with tables, subsidiary bosons**
- **Miscellaneous: batch modes, ROOT output, LHA formats, MCDB, nuclear PDF's,...**


27 years of CompHEP project in 2016

Primary publication: 1989



CompHEP general structure, SINP MSU preprint 91-9/213, 1991

- 29 -



Last stable version CompHEP 4.5.2 ,
download possible from <http://comphep.sinp.msu.ru>

Main objectives


- **Automation of tree level diagram calculations**
- **“Unification” of symbolic and numerical calculation, unweighted event generation for detector simulation - a full computational chain for collider physics**
- **Interfacing to other generators (partonic showering, hadronization, masses and mixings)**
- **Interfacing to NLO codes: cross section calculators, mass spectrum calculators**

Features

- **Generation of complete gauge invariant sets of tree-level Feynman diagrams**
- **Symbolic calculation of squared diagrams**
- **Generation of binary for numerical integration by Monte-Carlo method and calculation of cross sections and distributions**
- **Unweighted events generation**
- **Convenient format of built-in models. CompHEP can work with 0,1/2,1-spin particles, Majorana and Dirac spinors, 3- and 4-vertices with fields, derivatives of fields, functions of model parameters**
- **User-friendly interface: GUI for both symbolic and numerical parts, comprehensive built-in help (F1), batch scripts**
- **Generation of models by means of LanHEP (see <http://theory.sinp.msu.ru/~semenov/lanhep.html>)**

Distribution of citations: theory


CompHEP, NIM, A534 (2004) 250, 560 citations (period 2005 - june 2016)



Exotic BSM \Rightarrow scalar and vector leptoquarks, leptons and quarks of 4th generation, dileptons, mirror fermions, invisible H, little H, strong EW SB, color in the SB sector...

Distribution of citations: experimental analyses and simulations

CompHEP, NIM, A534 (2004) 250, 560 citations (period 2005 - june 2016)



**General approach of CompHEP, GRACE, 1990,
has been reproduced after 2000 and extended:**

Automatic event generation by ATLAS,CMS, etc.

(1) Sherpa + OpenLoops, FeynRules interface

**(2) Madgraph5 + Pythia6, MC@NLO,
MEPS@NLO, Mi@NLO, MadLoop,
FeynRules interface**

(3) Omega/Whizard, LO and NLO


Automatic check of models and hypotheses:

Checkmate, Delphes, Gambit

[more details E.Boos,M.D. Phys.Usp. 53 (2010) 1039]

Global fits

The signal strength and the signal strength error for various groups of Higgs boson production channels



Overall signal strength – all channels

$$1.00^{+0.14}_{-0.13} \left[\pm 0.09(\text{stat.})^{+0.08}_{-0.07}(\text{theo.}) \pm 0.07(\text{syst.}) \right]$$

Signal strength and exclusion contours in the SME (Standard Model Extension) parameter space

$$(1) \quad \mu_i = \frac{\left[\sum_{j=1}^{N_{\text{ch}}} \sigma_{j \rightarrow H} \text{Br}(H \rightarrow i) \right]_{\text{SME}}}{\left[\sum_{j=1}^{N_{\text{ch}}} \sigma_{j \rightarrow H} \text{Br}(H \rightarrow i) \right]_{\text{SM}}} \quad (2) \quad \mu_i = \frac{\left[\sum_{j=1}^{N_{\text{ch}}} \sigma_{j \rightarrow H(\text{off-shell}) \rightarrow i} \right]_{\text{SME}}}{\left[\sum_{j=1}^{N_{\text{ch}}} \sigma_{j \rightarrow H(\text{off-shell}) \rightarrow i} \right]_{\text{SM}}}$$

(1) signal strength in the production \times decay approximation

(narrow width approximation or infinitely small width approximation);

(2) signal strength for complete gauge invariant set

$$\hat{\mu}_i = \frac{N_{\text{obs},i} - N_{\text{backgr},i}}{N_{\text{signal},i}^{\text{SM}}}$$

- best fit of the signal strength for the number of experimentally observed signal events N_{OBS} , the number of background events N_{BACKGR} and the number of Standard Model events $N_{\text{SIGNAL}}^{\text{SM}}$;


$$\chi_{N_{\text{ch}}}^2 = \sum_{i=1}^{N_{\text{ch}}} \frac{(\mu_i - \hat{\mu}_i)^2}{\sigma_i^2}$$

MC - $\chi_{N_{\text{ch}}}^2$ distribution for the number of production channels N_{CH} ;


Beyond the infinitely small width approximation

In a number of channels the interference terms are not small (especially for $\gamma\gamma$, WW and ZZ exchange diagrams). Individual contributions of t-channel and subleading s-channel diagrams are usually small, but the number of such diagrams can be of the order of 100 (especially $\mu\mu\mu\mu$)


Example: $gg \rightarrow (W^*W^*, Z^*Z^*) \rightarrow \nu_\mu \nu_\mu \mu^+ \mu^-$




diagr.1




diagr.2




diagr.3



diagr.4



diagr.5



diagr.6

- **Up to which degree the SM Higgs boson is consistent with the available data? More than 200 production \otimes decay combinations and rearrangements are measured**
- **Structure of the couplings can be extracted correlating event rates from all channels**
- **Deviations from the SM are introduced in the form of effective operators O . Anomalous couplings C parametrize the deviations**

$$L_{eff}^{(6)} = \frac{1}{\Lambda^2} \sum_{k=V,F} C_{k\Phi} O_{k\Phi}$$


- **Global fit in the anomalous coupling space is performed combining all production channels**

E.Boos, V.Bunichev, M.D.,Y.Kurihara Phys.Rev.D 2014, Phys.Lett.B 2014


Uses signal strength definition (2) – complete gauge invariant sets

(c_V, c_F) parametrization.

c_V rescales the VVH, c_F rescales the FFH




$$c_F = 1 + C_{t\Phi} \cdot \frac{v^2}{\Lambda^2}$$



$$c_V = 1 + \frac{v^2}{2\Lambda^2} \cdot C_{\Phi}^{(1)}$$

$$c_G = c_F + \frac{6\pi}{\alpha_s} \cdot C_{\Phi G} \cdot \frac{v^2}{\Lambda^2}$$



$$c_\gamma = \frac{63c_F - 16c_V}{47} + \frac{9\pi}{4\alpha} \cdot (c_w^2 \cdot C_{\Phi B} + s_w^2 \cdot C_{\Phi W}) \cdot \frac{v^2}{\Lambda^2}$$

$$c_Z = (s_w^2 \cdot C_{\Phi B} + c_w^2 \cdot C_{\Phi W}) \cdot \frac{v^2}{\Lambda^2}$$

$$c_W = C_{\Phi W} \cdot \frac{v^2}{\Lambda^2}$$


the SM limit [$c_F=1, c_V=c_G=c_\gamma=1, c_W=0, c_Z=0$]

with the one-loop induced $H \rightarrow gg, H \rightarrow \gamma\gamma$ is clearly seen.

Effective triple vertices with the (c_F, c_V) parametrization

Triple vertices	Feynman rules
$\bar{t} \quad t \quad H$	$-\frac{M_t}{v} \cdot c_F$
$\bar{b} \quad b \quad H$	$-\frac{M_b}{v} \cdot c_F$
$\bar{\tau} \quad \tau \quad H$	$-\frac{M_\tau}{v} \cdot c_F$
$G_\mu \quad G_\nu \quad H$	$-\frac{2}{v} \cdot \frac{\alpha_s}{6\pi} \cdot c_G \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$A_\mu \quad A_\nu \quad H$	$-\frac{2}{v} \cdot \frac{4\alpha}{9\pi} \cdot c_\gamma \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$A_\mu \quad Z_\nu \quad H$	$+2 \cdot c_w \cdot s_w \cdot (C_{\Phi B} - C_{\Phi W}) \cdot \frac{v}{\Lambda^2} (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$Z_\mu \quad Z_\nu \quad H$	$+\frac{2}{v} \cdot [M_Z^2 \cdot c_V \cdot g^{\mu\nu} - c_Z \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)]$
$W_\mu^+ \quad W_\nu^- \quad H$	$+\frac{2}{v} \cdot [M_W^2 \cdot c_V \cdot g^{\mu\nu} - c_W \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)]$

Basic object: χ^2 measure in the anomalous coupling space.
Global fits to μ in (cF,cV) plane are performed. Dispersion matrix of the observables convoluted with vector differences between the observed and calculated μ values defines χ^2 . The minimum of χ^2 is found and 65%,90% and 99% best fit CL regions in the (cF,cV) space are defined by deviations from χ^2_{\min} less than 2.1,4.6 and 9.2, respectively.



New features of CompHEP v. 4.6 useful for generation of global fits

Implementation of external functions in the Constraints Model Table

Multiplication of selected squared diagrams on an external function

Table calculations and algebraic operations with tables — cross section/width vs parameters

**ROOT code generation to draw table functions
(3D surfaces or 2D contours)**

Generation of 3-DIM phase space distributions dependent on a model parameter

CompHEP Standard Model

The screenshot displays the CompHEP version 4.5.0rc6 interface, divided into four main panels:

- Variables:** A table listing physical constants and parameters. The selected row is 'EE' (Elementary charge).
- Particles:** A table listing particles with their properties. The selected row is 'tau-neutrino'.
- Constraints:** A table listing mathematical constraints. The selected row is 'CW'.
- Lagrangian:** A table listing Lagrangian terms. The selected row is 'P1'.

Variables Panel:

Name	Value	> Comment	<
EE	0.31345	Elementary charge (alpha=1/127.9, on-shell, MZ)	
GG	1.21358	Strong coupling constant (Z pnt, alp=0.1172pm0)	
SW	0.48076	sin of the Weinberg angle (MZ point -> MW=79.9)	
s12	0.2229	Parameter of C-K-M matrix (PDG2002)	
s23	0.0412	Parameter of C-K-M matrix (PDG2002)	
s13	0.0036	Parameter of C-K-M matrix (PDG2002)	
MZ	91.1876	mass of Z boson	
wZ	2.43631	width of Z boson	
wW	2.02798	width of W boson	
Mm	0.10566	mass of muon	
Mtau	1.77699	mass of tau-lepton	
Mc	1.65	mass of c-quark	
Ms	0.117	mass of s-quark	
Mtop	174.3	mass of t-quark	
wtop	1.54688	width of t-quark	
Mb	4.85	mass of b-quark	
MH	115	mass of Higgs	
wH	0.0061744	width of Higgs	

Particles Panel:

Full name	P	aP	2*spin	mass	width	color	aux	>	LaTeX(A)	<
gluon	G	G	2	0	0	8	G	G	G	G
photon	A	A	2	0	0	1	G	A	A	A
Z boson	Z	Z	2	MZ	wZ	1	G	Z	Z	Z
W boson	W+	W-	2	MW	wW	1	G	W+	W^	W^
neutrino	ne	Ne	1	0	0	1	L	nu^e	nu^e	nu^e
electron	e	E	1	0	0	1		e	e	e
mu-neutrino	nm	Nm	1	0	0	1	L	nu^mu	nu^mu	nu^mu
muon	m	M	1	Mm	0	1		mu	mu	mu
tau-neutrino	nl	Nl	1	0	0	1	L	nu^tau	nu^tau	nu^tau
tau-lepton	l	L	1	Mtau	0	1		tau	tau	tau
u-quark	u	U	1	0	0	3		u	u	u
d-quark	d	D	1	0	0	3		d	d	d
c-quark	c	C	1	Mc	0	3		c	c	c
s-quark	s	S	1	Ms	0	3		s	s	s
t-quark	t	T	1	Mtop	wtop	3		t	t	t
b-quark	b	B	1	Mb	0	3		b	b	b
Higgs	H	H	0	MH	wH	1		H	H	H

Constraints Panel:

Name	> Expression	<
CW	sqrt(1-SW^2)	
c12	sqrt(1-s12^2)	
c23	sqrt(1-s23^2)	
c13	sqrt(1-s13^2)	
Vud	c12*c13	
Vus	s12*c13	
Vub	s13	
Vcd	s12*c23-c12*s23*s13	
Vcs	c12*c23-s12*s23*s13	
Vcb	s23*c13	
Vtd	s12*s23-c12*c23*s13	
Vts	c12*s23-s12*c23*s13	
Vtb	c23*c13	
MW	MZ*CW	

Lagrangian Panel:

P1	P2	P3	P4	> Factor	<
P1	b	W+		EE*Sqrt2*Vcb/(4*SW)	G(m)
C	b	W+.f		i*EE*Sqrt2*Vcb/(4*MW*SW)	Mb*
C	c	A		EE/3	G(m)
C	c	G		GG	G(m)
C	c	H		EE*Mc/(2*MW*SW)	1
C	c	Z		EE/(12*CW*SW)	3-
C	c	Z.f		i*EE*Mc/(2*MW*SW)	G5
C	d	W+		EE*Sqrt2*Vcd/(4*SW)	G(m)
C	d	W+.f		i*EE*Mc*Sqrt2*Vcd/(4*MW*SW)	1-
C	s	W+		EE*Sqrt2*Vcs/(4*SW)	G(m)
C	s	W+.f		i*EE*Sqrt2*Vcs/(4*MW*SW)	Ms*
D	c	W-		EE*Sqrt2*Vcd/(4*SW)	G(m)
D	c	W-.f		i*EE*Mc*Sqrt2*Vcd/(4*MW*SW)	1+
D	d	A		EE/3	G(m)
D	d	G		GG	G(m)
D	d	Z		EE/(12*CW*SW)	2*S
D	t	W-		EE*Sqrt2*Vtd/(4*SW)	G(m)
D	t	W-.f		i*EE*Mtop*Sqrt2*Vtd/(4*MW*SW)	1+
D	u	W-		EE*Sqrt2*Vud/(4*SW)	G(m)
E	e	A		EE	G(m)
E	e	Z		EE/(4*CW*SW)	1-

Using external functions in the Constraints Model Table

Any model parameter and vertex form-factor may be represented in the form of «c»-function that depends on other model parameters and on 4-momenta of particles

```
CompHEP version 4.6
Constraints 17
Clr Rest Del Size
Name |> Expression
coeff |coeff1(MR,sint)
wH |width1(MR,sint)
wR |width2(MR,sint)
c |width3(MR,sint)
b |b1*c
yt |myfunc2(Mtop)
yW |myfunc2(MW)
loopt |myfunc3(yt,1)
loopW |myfunc3(yW,2)
ImIt |myfunc4(yt,1)
ImIW |myfunc4(yW,2)
RFF |1-(b1*cos+ b*sint-sint/v)
HFF |1-(b1*sint+ b*cos+ cost/v)
Ranom |b1*cos- b*sint
Hanom |b1*sint+ b*cos
RGG |17*Ranom+loopt*(-RFF)
ImRGG |ImIt*(-RFF)
HGG |17*Hanom+loopt*(-HFF)
ImHGG |ImIt*(-HFF)
RAA |1-11/(3)*Ranom+(loopW+8/(3)*loopt)*(-RFF)
ImRAA |1(ImIW+8/(3)*ImIt)*(-RFF)
HAA |1-11/(3)*Hanom+(loopW+8/(3)*loopt)*(-HFF)
ImHAA |1(ImIW+8/(3)*ImIt)*(-HFF)
F1 F2 Top Bottom GoTo Find Zoom ErrMes
```

```
myfunc.c
double myfunc3 (double ym, double keyp)
{
    double result, Fym, as, sqr, logs;

    as = asin(1./sqrt(fabs(ym)));
    sqr = sqrt(fabs(1.-ym));
    logs = log((1.+sqr)/(1.-sqr));

    if(ym >= 1.0) Fym = as*as;
    else Fym = -0.25*(logs*logs-9.869587728);


    if(keyp < 1.5) result = ym*(1.+(1.-ym)*Fym);
    else result = -(2. + 3.*ym + 3.*ym*(2.-ym)*Fym);

    return result;
}
```

Multiplication of selected squared diagrams on an external function



```

CompHEP version 4.6
Constraints 17
Clr Rest Del Size
Name |> Expression
coeff |coeff1(MR,sint)
wH |width1(MR,sint)
wR |width2(MR,sint)
c |width3(MR,sint)
b |b1*c
yt |myfunc2(Mtop)
yW |myfunc2(MW)
loopt |myfunc3(yt,1)
loopW |myfunc3(yW,2)
ImIt |myfunc4(yt,1)
ImIW |myfunc4(yW,2)
RFF |-(b1*cost-b*sint-sint/v)
HFF |-(b1*sint+b*cost+cost/v)
Ranom |b1*cost-b*sint
Hanom |b1*sint+b*cost
RGG |7*Ranom+loopt*(-RFF)
ImRGG |ImIt*(-RFF)
HGG |7*Hanom+loopt*(-HFF)
ImHGG |ImIt*(-HFF)
RAA |-11/(3)*Ranom+(loopW+8/(3)*loopt)*(-RFF)
ImRAA |ImIW+8/(3)*ImIt*(-RFF)
HAA |-11/(3)*Hanom+(loopW+8/(3)*loopt)*(-HFF)
ImHAA |ImIW+8/(3)*ImIt*(-HFF)
F1 F2 Top Bottom GoTo Find Zoom ErrMes
    
```



One can mark some squared diagrams in GUI mode, these diagrams are then multiplied by the function «coeff», where «coeff» is an external "c" -function or two-dimensional table

Algebraic operations with tables –cross section/width vs parameters



CompHEP version 4.6

```

: G,G -> A,A
session: 1(begin)
    
```

Numerical Session

*
 Itmx = 5
 nCall = 98568

Distributions
 Start integration
 Clear statistic

*
 Define variables
 Display 1d distributions
 Combine 1d distributions
 Sum 1d distributions

Sum 1d tables
 Plot 1d table
 Construct 2d distribution

Sum 2d tables
 Add constant to table terms
 Multiply 2d tables
 Multiply 2d table to koeff.
 Divide 2d tables
 Rescale x variable
 Rescale y variable

Plot 3d surface
 Plot 2d contour

Calculate hi^2
 Sum hi^2 tables
 Plot delta hi^2 contour
 Erase data

$\Gamma_{H \rightarrow 2X} + \Gamma_{H \rightarrow WW^*, ZZ^*}$

=

lp F2-Man F4-Diagrams F5-Squared Diagrams F6-Results F9-Quit

resulting tables can be used as external functions in the Constraints Model Table

```

CompHEP version 4.6
Constraints 18
Clr Rest Del Size
Name |> Expression
coeff |coeff1(MR,sint)
wH |width1(MR,sint)
wR |width2(MR,sint)
c |width3(MR,sint)
b |b1*c
yt |myfunc2(Mtop)
yW |myfunc2(MW)
loopt |myfunc3(yt,1)
loopW |myfunc3(yW,2)
ImIt |myfunc4(yt,1)
ImIW |myfunc4(yW,2)
RFF |-(b1*cost-b*sint-sint/v)
HFF |-(b1*sint+b*cost+cost/v)
Ranom |b1*cost-b*sint
Hanom |b1*sint+b*cost
RGG |7*Ranom+loopt*(-RFF)
ImRGG |ImIt*(-RFF)
HGG |7*Hanom+loopW*(-HFF)
ImHGG |ImIW*(-HFF)
RAA |-11/(3)*Ranom+(loopW+8/(3)*loopt)*(-RFF)
ImRAA | (ImIW+8/(3)*ImIt)*(-RFF)
HAA |-11/(3)*Hanom+(loopW+8/(3)*loopW)*(-HFF)
ImHAA | (ImIW+8/(3)*ImIW)*(-HFF)
F1-F2 Top Bottom GoTo Find Zoom ErrMes


```

width1.txt x


1.000000E+02	-7.071070E-01	9.265819E-04
1.000000E+02	-6.788227E-01	1.013980E-03
1.000000E+02	-6.505384E-01	1.099670E-03
1.000000E+02	-6.222542E-01	1.183458E-03
1.000000E+02	-5.939699E-01	1.265209E-03
1.000000E+02	-5.656856E-01	1.344767E-03
1.000000E+02	-5.374013E-01	1.422014E-03
1.000000E+02	-5.091170E-01	1.496865E-03
1.000000E+02	-4.808328E-01	1.569201E-03
1.000000E+02	-4.525485E-01	1.638916E-03
1.000000E+02	-4.242642E-01	1.705987E-03
1.000000E+02	-3.959799E-01	1.770256E-03
1.000000E+02	-3.676956E-01	1.831733E-03
1.000000E+02	-3.394114E-01	1.890312E-03
1.000000E+02	-3.111271E-01	1.945908E-03
1.000000E+02	-2.828428E-01	1.008526E-02

ROOT code generation to draw table functions (3D surfaces or 2D contours)

$gg \rightarrow \gamma\gamma$ (LHC, $\sqrt{s}=8$ TeV, $m_h=125$ GeV, $\Lambda_r=3$ TeV, $c=c_{\max}$)



$r \rightarrow \gamma\gamma$ tag (LHC, $\sqrt{s}=13$ TeV, $m_t=175$ GeV, $c=c_{\max}$)



Numerical Session

* Itmx = 5
nCall = 98568

Distributions

* Define variables
Display 1d distributions
Combine 1d distributions
Sum 1d distributions

Sum 1d tables
Plot 1d table
Construct 2d distribution

Sum 2d tables
Add constant to table terms
Multiply 2d tables
Multiply 2d table to koeff.
Divide 2d tables
Rescale x variable
Rescale y variable

Plot 3d surface

Plot 2d contour

Calculate hi^2
Sum hi^2 tables
Plot delta hi^2 contour
Erase data


2-Man F4-Diagrams F5-Squared Diagrams F6-Results F9-Quit

ROOT code generation for 3D phase space distributions dependent on a BSM model parameter

CompHEP version 4.6

(sub
Monte

$u\bar{d} \rightarrow t\bar{b}$, t-quark p_t



$d\sigma/dp_t$, fb/GeV

p_t

f_{RT}

Numerical Session

*
Itmx = 5
nCall = 98568

Distributions

*
Define variables
Display 1d distributions
Combine 1d distributions
Sum 1d distributions





Sum 1d tables
Plot 1d table
Construct 2d distribution

Sum 2d tables
Add constant to table terms
Multiply 2d tables
Multiply 2d table to coeff.
Divide 2d tables
Rescale x variable
Rescale y variable
Plot 3d surface
Plot 2d contour

Calculate h_i^2
Sum h_i^2 tables
Plot Δh_i^2 contour
Erase data

F1-Help F2-Man F4-Diagrams F5-Squared Diagrams F6-Results F9-Quit

**LHC
2014
ATLAS
+CMS
combined**



Technical problems of evaluations with higher dim operators of BSM

- **several anomalous couplings (AC) from different effective operators contribute to $|M|^2$**
- **different AC contribute to the decay widths of unstable particles**
- **from other side, contributions of individual AC are used for event samples in experimental searches**

Separation of congenerous contributions (e.g. $1/\Lambda^2$ leading terms) in the event samples is of interest

Subsidiary bosons for BSM evaluations

New Physics (NP) contributions to the SM vertex

$$\Gamma_\mu = \Gamma_\mu^{\text{SM}} + \Gamma_\mu^{\text{NP}_1} + \Gamma_\mu^{\text{NP}_2} + \dots$$

Example: anomalous Wtb vertex

$$\mathcal{L}_{\text{Wtb}} = \frac{g}{\sqrt{2}} \bar{b} \gamma^\mu (f_V^L P_L + f_V^R P_R) t W_\mu^- + \frac{g}{\sqrt{2}} \bar{b} \frac{\sigma^{\mu\nu}}{m_W} (f_T^L P_L + f_T^R P_R) t W_{\mu\nu}^- + h.c.$$


W boson SM $\frac{g}{2\sqrt{2}} f_V^L \gamma^\mu (1 - \gamma_5)$

W boson subsidiary 1 $\frac{g}{2\sqrt{2}} f_V^R \gamma^\mu (1 + \gamma_5)$

W boson subsidiary 2 $\frac{g}{2m_W\sqrt{2}} f_T^L \sigma^{\mu\nu} q_\nu (1 + \gamma_5)$

W boson subsidiary 3 $\frac{g}{2m_W\sqrt{2}} f_T^R \sigma^{\mu\nu} q_\nu (1 - \gamma_5)$

Boos, Bunichev, Dudko, Perfilov, arXiv:1512.00826, arXiv:1607.00505



Diagrams (2),(3),(4) with subsidiary bosons for $qq \rightarrow bb \mu \nu_\mu$
Squared amplitude with 'production' P_1, P_2 and 'decay' D_1, D_2

$$\begin{aligned}
 |M|^2 &\sim \frac{1}{\Gamma} [(f_V^L)^2 P_1 + (f_V^R)^2 P_2] \times [(f_V^L)^2 D_1 + (f_V^R)^2 D_2] \\
 &\sim \frac{1}{\Gamma} [(f_V^L)^4 P_1 D_1 + (f_V^L)^2 (f_V^R)^2 P_1 D_2 + (f_V^L)^2 (f_V^R)^2 P_2 D_1 + (f_V^R)^4 P_2 D_2]
 \end{aligned}$$

Three sets of event samples for simulation when $f_{LV}=f_{RV}=1, f_{LT}=f_{RT}=0$

$$(f_V^L f_V^R 00) \Leftrightarrow (f_V^L)^4 \otimes (1000) \oplus (f_V^L)^2 (f_V^R)^2 \otimes (1100)_{\text{sub}} \oplus (f_V^R)^4 \otimes (0100)_{\text{sub}}$$

Physics Analysis Summary CMS-PAS-TOP-14-007. Baesian Neural Network Discriminant (BNN)

CMS preliminary, $\sqrt{s} = 7$ TeV, $L = 5.0$ fb $^{-1}$






Figure 7: Data and model comparison of BNN aWtb discriminant for the (f_V^L, f_T^L) scenario. The BNN aWtb was trained to separate possible events with left tensor coupling in the Wtb interaction and SM events. The hashed band corresponds to the systematic uncertainty.

Physics Analysis Summary CMS-PAS-TOP-14-007

CMS preliminary, $\sqrt{s} = 7 \text{ TeV}, L = 5.0 \text{ fb}^{-1}$



CMS preliminary, $\sqrt{s} = 7 \text{ TeV}, L = 5.0 \text{ fb}^{-1}$



Important features improved


- **Batch system. Symbolical and numerical batch calculations in PBS/LSF**
- **Output event format respecting Les Houches agreements (LHEF with HepML header), convention LHAPDF, SUSY LHA format, BSM LHA format)**
- **Interfaces to PYTHIA/HERWIG and other**
- **Monte Carlo events data base (MCDB, see Comput. Phys. Commun.178(2008)222, hep-ph/0703287)**
- **Nuclear PDF's (Phys.Rev.C92(2015)044901, hep-ph/0703287)**

Summary

- **CompHEP developments in 2010-2016 have been motivated mainly by experimental analyses of CMS and D0 collaborations. Tools for identification of the Higgs boson and the top quark have been developed.**
- **External functions, operations with cross section/Br tables, generation of combined fits and implementation of subsidiary fields are introduced to work in the BSM multiparameter space.**
- **Visualization, batch modes and interfaces significantly improved.**

Backup slides

Modern Monte-Carlo Chain



General information and references

- CompHEP collaboration: E. Boos, V. Bunichev, M. Dubinin, L. Dudko, V. Ilyin, A. Kryukov, V. Edneral, V. Savrin (Moscow State), A. Semenov (JINR, Dubna), A. Sherstnev
- CompHEP homepage: <http://comphep.sinp.msu.ru>
- References:
 - CompHEP 4.5 Status Report. E.Boos et al. arXiv:0901.4757
 - CompHEP: E. Boos et al., Nucl.Inst.Meth. A534:250 (2004) [hep-ph/0403123]
 - LanHEP: A. Semenov, Nucl.Inst.Meth. A393:293 (1997) [hep-ph/0403123]; 0805.0555 (hep-ph)
 - CompHEP-Interfaces: A.Belyaev et al., hep-ph/0101232

Les Houches Agreements

There are many MC generators with their own advantages and application areas. Often we are forced to use several generators for reliable calculations:

Problems:

- Interfacing some MC codes (ME and SH generators): Les Houches Accord 1, Les Houches Event format
- Les Houches Accord 2: uniform interface to different PDF sets (LHAPDF package)
- Les Houches Accord 3: Interfacing SUSY codes to MC generators for parameters, spectrum, decays (SPA).
- BSM Les Houches Accord: fixing of parameter record for BSM
- Matching ME (LO/NLO) and SR(NL): CKKW, MC@NLO, Mrenna-Richardson, MLM, ...

Batch system in CompHEP

**Both symbolic & numerical parts of the package have batch scripts:
symb_batch.pl and num_batch.pl (in Perl)**

Useful in the cases

- **Computations of many (of the order of 100) subprocesses for LHC analyses**
- **Remote calculations:** GUI not convenient
- **Support of parallel calculations:** very helpful for multi-CPU machines/computer clusters (pbs/lsf is available; grid in progress)

Symbolical batch: pp->m,Nm,b,B,H+ with t->b,H+ and T->m,Nm,B MSSM, tb=0.5, MH+=150GeV (H+->t*b->2f+bB dominates)

27

- Prepare `process.dat` following toy example: all points well documented
- `./symb_batch.pl -show diag` (to exclude several sub-leading diagrams) diagrams in 9 subprocesses (54 sqr. diag.) (15 G,G->m,Nm,b,B,H+ diagrams)
- `./symb_batch.pl -mp 2` calculate faster (2 times if you have 2*CPU machine)

```
#####
# Data file for symb_script.pl
# For the symb_batch script version 1.0
#####

# You have to set the model number, which you are going to
# The model number corresponds to the string number of the
# in the CompHEP model menu in the GUI mode..
model number: 6

# Beam names can be taken from a table of beams.
# (see CompHEP in the GUI regime). Energy unit is GeV
beam 1: p
beam 2: p
beam energy 1: 7000.0
beam energy 2: 7000.0


# This string defines the final state of your process. Mode
# particles and composite particles (see the corresponding
# can be used..
final state: m,Nm,b,B,H+

# If you'd like to exclude feynman diagrams with some model
# particles (in propagators!), enter the particles here]
exclude diagrams with: h,H,H3,u,d,c,s,A,Z

# If you'd like to keep feynman diagrams with some model
# particles (in propagators!), enter the particles here
# Examples:
#keep diagrams with: t,b,Z,A
keep diagrams with:

# If you enter no, s_comphеп generates diagrams and does no
# do symbolic calculations.
make symbolic calculations(yes/no): yes

# If you enter no, comphep calculates all squared diagrams,.
# but n_comphеп will not be created.
make n_comphеп generator(yes/no): yes
```



```
[note]$ ./symb_batch.pl -show stat
Diagram statistics: total = 54, calculated = 44, deleted = 0
[note]$ Old n_comphеп is deleted!
End of CompHEP symbolical session.
*** n_comphеп creation details have been written to symb_batch.log
```

Numerical batch: pp->m,Nm,b,B,H+ in MSSM

- Prepare **batch.dat**: customize first process via GUI and execute **./num_batch.pl**
- Customize differences in other subprocesses (if needed) via GUI and execute **./num_batch.pl -add -proc ...** for the necessary subprocesses
- Start numerical calculations with **./num_batch.pl -run ...**

```
#Subprocess 1 (u,U -> m,Nm,b,B,H+)
#Session_number 1
#Model_number 6
#Initial_state
  SQR(T(S)) 1.400000E+04
  Rapidity(c.m.s) 0.000000E+00
  StrFun1: PDF:cteq6l1(proton)
  StrFun2: PDF:cteq6l1(proton)

#Physical_Parameters
  EE = 3.1223000000000000E-01
  SW = 4.7300000000000000E-01
  MZ = 9.1188400000000000E+01
  Mtop = 1.7500000000000000E+02
  Mb = 4.6200000000000000E+00
  wtop = 1.7524000000000000E+00
  ww = 2.0889500000000000E+00
  mu = 1.0000000000000000E+03
  MG2 = 2.0000000000000000E+02
  MG3 = 3.0000000000000000E+02
  Mq3 = 1.0000000000000000E+03
  Mu3 = 1.0000000000000000E+03
  Md3 = 1.0000000000000000E+03
  Atop = 0.0000000000000000E+00
  Ab = 0.0000000000000000E+00
  MH3 = 1.3416000000000000E+02
  tb = 5.0000000000000000E-01
  GG = 1.216002374681738E+00
```

```
#Width_scheme 0

#Kinematical_scheme
12 -> 57 , 346
57 -> 5 , 7
346 -> 6 , 34
34 -> 3 , 4

#Cuts.
```

```
[note]$ ./num_batch.pl --show cs
List of available subprocesses:
```

```
Subprocess 1 (u,U -> m,Nm,b,B,H+): cross section [pb] = 6.2925e-01 +/- 1.30e-03 ( 2.06e-01 % )
Subprocess 2 (d,D -> m,Nm,b,B,H+): cross section [pb] = 3.8960e-01 +/- 8.15e-04 ( 2.09e-01 % )
Subprocess 3 (U,u -> m,Nm,b,B,H+): cross section [pb] = 6.2781e-01 +/- 1.55e-03 ( 2.47e-01 % )
Subprocess 4 (D,d -> m,Nm,b,B,H+): cross section [pb] = 3.8906e-01 +/- 9.31e-04 ( 2.39e-01 % )
Subprocess 5 (s,S -> m,Nm,b,B,H+): cross section [pb] = 6.6678e-02 +/- 1.43e-04 ( 2.14e-01 % )
Subprocess 6 (c,C -> m,Nm,b,B,H+): cross section [pb] = 3.0779e-02 +/- 6.58e-05 ( 2.14e-01 % )
Subprocess 7 (S,s -> m,Nm,b,B,H+): cross section [pb] = 6.6678e-02 +/- 1.43e-04 ( 2.14e-01 % )
Subprocess 8 (C,c -> m,Nm,b,B,H+): cross section [pb] = 3.0779e-02 +/- 6.58e-05 ( 2.14e-01 % )
Subprocess 9 (G,G -> m,Nm,b,B,H+): cross section [pb] = 1.4684e+01 +/- 3.59e-02 ( 2.44e-01 % )
```

```
Total CS [pb] = 1.6914e+01 +/- 3.60e-02 ( 2.13e-01 % )
```

```
#Regularization.
```

```
*** Table ***
```

```
Regularization
Momentum |> Mass <|> Width <|> Power|
57         |Mtop  |wtop  |2.....
34         |MW    |ww    |2.....
346        |Mtop  |wtop  |2.....
```

```
#QCD Lambda6 = 1.652000E-01 Scale = 175
#Vegas_calls 41472x5
#Vegas_integral 9.16788703338995469E+13 3.46369076228:
#Distributions.
```

```
*** Table ***
```

```
Distributions
```

```
Parameter |> Min bound <|> Max bound <|> Rest Frame
```

```
=====
#Events 500 1 0.200000 2.000000 10000
#Random FA98C8AA370E
```

```
#VEGAS Grid Vegas grid: dim=12 size=50
```

Sector by sector extension of the SM by dimension 5 and 6 effective operators

W.Buchmuller, D.Wyler, Nucl.Phys. B268 (1986) 621

Recent two-parametric global fits – nonlinear chiral realization of the SM gauge symmetry (alternative)

J.R. Espinosa, C. Grojean, M. Muhlleitner, M. Trott, JHEP 1205, 097 (2012)

(arXiv:1202.3697 [hep-ph]), JHEP 1212, 045 (2012) (arXiv:1207.1717 [hep-ph])

- *scalar-gauge boson sector*

$$O_{\Phi G} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})G_{\mu\nu}^a G^{a\mu\nu}$$

$$O_{\Phi B} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})B_{\mu\nu} B^{\mu\nu}$$

$$O_{\Phi W} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})W_{\mu\nu}^i W^{i\mu\nu}$$

$$O_{\Phi}^{(1)} = (\Phi^\dagger\Phi - \frac{v^2}{2})D_\mu\Phi^\dagger D^\mu\Phi$$

$$O_{\Phi G} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})G_{\mu\nu}^a \bar{G}^{a\mu\nu}$$

$$O_{\Phi B} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})B_{\mu\nu} \bar{B}^{\mu\nu}$$

$$O_{\Phi W} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})W_{\mu\nu}^i \bar{W}^{i\mu\nu}$$

- *scalar-fermion sector*

$$O_{t\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(\bar{Q}_L\Phi^c t_R)$$

$$O_{b\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(\bar{Q}_L\Phi b_R)$$

$$O_{\tau\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(\bar{L}_L\Phi\tau_R)$$

$$\bar{F}_{\mu\nu} = \epsilon_{\mu\nu\gamma\delta}F_{\gamma\delta}.$$

Effective triple vertices in the Buchmueller-Wyler basis (LanHEP calculation). Effective couplings C (Wilson coefficients) are multiplicative factors in front of O_{ij}

Effective operators	Triple vertices	Feynman rules
$O_{t\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(-\lambda_t)(\bar{Q}_L\Phi^c t_R)$	$\bar{t} \quad t \quad H$	$-M_t \cdot \frac{v}{\Lambda^2} \cdot C_{t\Phi}$
$O_{b\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(-\lambda_b)(\bar{Q}_L\Phi b_R)$	$\bar{b} \quad b \quad H$	$-M_b \cdot \frac{v}{\Lambda^2} \cdot C_{b\Phi}$
$O_{\tau\Phi} = (\Phi^\dagger\Phi - \frac{v^2}{2})(-\lambda_\tau)(\bar{L}_L\Phi\tau_R)$	$\bar{\tau} \quad \tau \quad H$	$-M_\tau \cdot \frac{v}{\Lambda^2} \cdot C_{\tau\Phi}$
$O_{\Phi G} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})G_{\mu\nu}^a G^{a\mu\nu}$	$G_\mu \quad G_\nu \quad H$	$-2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi G} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$O_{\Phi B} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})B_{\mu\nu} B^{\mu\nu}$	$A_\mu \quad A_\nu \quad H$ $A_\mu \quad Z_\nu \quad H$ $Z_\mu \quad Z_\nu \quad H$	$-2 \cdot c_W^2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi B} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$ $+2 \cdot c_W \cdot s_W \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi B} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$ $-2 \cdot s_W^2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi B} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$O_{\Phi W} = \frac{1}{2}(\Phi^\dagger\Phi - \frac{v^2}{2})W_{\mu\nu}^i W^{i\mu\nu}$	$A_\mu \quad A_\nu \quad H$ $A_\mu \quad Z_\nu \quad H$ $Z_\mu \quad Z_\nu \quad H$ $W_\mu^+ \quad W_\nu^- \quad H$	$-2 \cdot s_W^2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi W} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$ $-2 \cdot c_W \cdot s_W \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi W} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$ $-2 \cdot c_W^2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi W} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$ $-2 \cdot \frac{v}{\Lambda^2} \cdot C_{\Phi W} \cdot (g^{\mu\nu} p_1 p_2 - p_1^\nu p_2^\mu)$
$O_\Phi^{(1)} = (\Phi^\dagger\Phi - \frac{v^2}{2})D_\mu\Phi^\dagger D^\mu\Phi$	$W_\mu^+ \quad W_\nu^- \quad H$ $Z_\mu \quad Z_\nu \quad H$	$M_W^2 \cdot \frac{v}{\Lambda^2} \cdot C_\Phi^{(1)} \cdot g^{\mu\nu}$ $M_Z^2 \cdot \frac{v}{\Lambda^2} \cdot C_\Phi^{(1)} \cdot g^{\mu\nu}$